# Lunar Volcanism

Angel Garlant Mentor: Dr. Lisa Gaddis

## Pyroclastic Deposits



- **™** Volcanic material
- CR Low albedo
- Smooth, rough, blocky
- - **Atlas**
  - Alphonsus
  - 3 J. Herschel
- **™** Volcanic source vents
  - S Floor fractures
  - **C**raters

## Volcanism

#### 03


- **Strombolian** 
  - Typically low silica content
  - **3** Bubbles
- **W** Vulcanian
  - Higher presence of silica
  - **3** Caprock
  - Intermittent

## Methods




- **Georeferencing** 
  - **S** ArcGIS
- **Crater Measurements** 
  - **S** ArcGIS
  - **US** ISIS
  - **G** Excel
- Modeling
  - **S** Excel

# Magma




## Ejection





# Trajectory

CS



### ArcGIS

#### CS

- ArcCatalog
  - Assign coordinate system to images
- ArcMap
  - Import images
  - Create tie points to relocate images to a different point
- Allows for easier viewing

# ISIS



- Craters are measured
- Screenshots are taken to keep track of which craters have been measured
- Can also be used to create a mosaic




### Excel

#### CF

- Used for entering data and doing computations
- Crater measurements are entered
  - Average diameter
  - **S** Rim height
  - Use ArcMap to locate coordinates of crater
- - Initial conditions (pressure, gas content, etc.)
  - **S** Range of pyroclasts


## J Herschel Crater






# Major Deposit







## Characteristics

03

₹ 70% olivine

≈ 30% pyroxene

№ 165 km diameter

Area: 60 km by 35 km

## Goals



- Revelop an isopach map
- Use model to run a simulation
- Revelop a general model
  - Are pyroclastic deposits from different areas erupted by different means
  - 13 How do they differ

# Questions?

03

